

TABLE 1. Measures related to calculating the number of women treated in health facilities for complications of induced abortion, by geographic region, Ethiopia, 2008

Region	No. of women treated in health facilities for abortion complications			No. of live births	No. of late miscarriages§
	All*	Miscarriage†	Induced abortion‡		
All	57,964	5,357	52,607	2,964,323	101,083
Tigray	3,338	337	3,001	163,444	5,573
Amhara	9,242	759	8,483	641,434	21,873
Oromiya	20,310	1,724	18,586	1,185,400	40,422
SNNP	10,896	773	10,123	618,066	21,076
Addis Ababa	7,233	1,077	6,156	40,564	1,383
4 rural regions**	3,274	520	2,754	297,683	10,151
2 urban regions††	3,671	167	3,504	17,732	605

*Includes both spontaneous and induced abortions. †Calculation assumes that the proportion of women with miscarriages who obtain treatment is the same as the proportion who deliver in facilities. ‡The total number treated for any abortion complication minus the number treated for complications of miscarriages. §Miscarriages at 13–21 weeks' gestation, calculated as 3.41% of all live births. **Includes Affar, Somali, Benishangul-Gumuz and Gambela. ††Includes Harari and Dire Dawa. Note: SNNP=Southern Nations, Nationalities and Peo-